


# Gastrointestinal System Clinical Examination


---



# CHALLENGE




# Anatomical areas




Abdomen - four quadrants

Copyright © 2003 by Lippincott Williams & Wilkins, *Instructor's Resource CD-ROM to accompany Bates' Guide To Physical Examination And History Taking*, 8th edition



# Order of GI tract examination




# ORGANS & FUNCTIONS

**Liver**  
Bile acids  
(aid lipid digestion)  
First-pass organ of  
ingested metabolites

**Duodenum**  
Neutralization  
Protein, lipid,  
carbohydrate digestion  
Absorption:  
Nutrients  
Electrolytes and  
metal ions  
Water

**Ileum**  
Absorption:  
Bile acids  
Vitamin B12  
Role in immunity



Churn and mix  
Protein digestion  
Acid production:  
Aids protein digestion  
Antimicrobial  
Intrinsic factor  
Iron reduction

**Pancreas**  
Digestive enzymes


**Jejunum**  
Digestion  
Absorption:  
Electrolyte-rich fluid

**Colon**  
Absorption:  
Water  
Electrolytes  
Bacterial metabolism  
Fatty acid metabolism

# Abdominal pain



1. VISCERAL
2. PARIETAL
3. SUPERFICIAL
4. REFERRED



Sources of abdominal pains

Copyright © 2003 by Lippincott Williams & Wilkins. *Instructor's Resource CD-ROM to Accompany Bates' Guide To Physical Examination And History Taking*, 8th edition

# Visceral pain

---

1. Spastic- induced by spasm of a hollow viscera, sudden, short, well localized, comes in cramps, relieved by thermoprocudures or spasmolytics (gall bladder, kidney, bowel, stomach)
2. Distensive – induced by distention by gases, feces, food, bile, has gradual onset, long standing, permanent, poorely localized (meteorism, hyposecretic syndrome)
3. Vascular (intestinal angina) – induced by ischemia, extremely severe (mesenteric thrombosis, spasm, arterial embolism)

1. **Parietal pain** – arises from impulses in the parietal peritoneum, well localized, accentuated by pressure, coughing, sneezing (peritonitis as a result of inflammation of an organ or perforation)

2. **Superficial pain** – abdominal wall pain (skin, nerves, muscles) it is sharp, constant and superficial, aggravated by contraction of abdominal musculature

3. **Referred pain** – is radiated from the affected organ (the area of reference has the same central pathways for afferent neurons ) Ex. Inferior myocardial infarction is felt in the epigastric area



# Characteristics of pain

- Location
- Onset
- Character described by adjectives—sharp/dull, Burning/ tingling, boring/stabbing, crushing/tugging.
- Radiation
- Associated symptoms
- Timing Since onset (episodic duration and frequency of attacks, the evolution)
- Aggravating and relieving factors (food or specific activities, postures or some medication)
- Severity subjective variation by day or night, week or month




# Extracardiac causes of chest pain

## Common Causes of Chest Pain

- Aortic → Aortic dissection, Aortic aneurism
- Esophageal/GI → Esophagitis, Esop. Spasm , esophageal tear  
Pancreatitis, Biliary /GB disease , GERD, Peptic Ulcer
- Lungs & Pleura → Bronchospasm, PE, Pneumonia ,TB,  
Trachitis, Pleuritis, PneumThorax , Malignancy  
, Asthma.
- Musculo-Skeletal → Ost. Arthritis, Rib#, I. Costal Muscle injury,  
Costochondritis, Cerv. Disc Disease
- Neurological → Prolapsed disc, Herpez Zoster,  
Thoracic Outlet Syndrome
- Psychological/  
others → Panic Attack/Anxiety Disorders ,  
Cocaine abuse

Figure 3: Differential Diagnoses According to Localization of Abdominal Pain produced by Shaza Karrar



# Character of pain

- ❑ **Gastric pain** could be dull, intense, “burning”;
  - Localized in epigastria; irradiating to median line
  - Associated with vegetative manifestations – nausea, transpiration
- ❑ **Intestinal pain** – has a colic character – pain periods alternating with periods of leisure; **colonic** pain – non-localized, in the whole abdominal cavity; **rectal** pain – in anal region, spreading to sacral region
- ❑ **Hepatic** pain – in right hypochondria
- ❑ **Gallbladder pain** – in epigastria, irradiate in right hypochondria, to right scapula.
- ❑ **Pancreatic pain** – in left hypochondria, epigastria, right hypochondria, like a “belt”.

# Timing

- ❖ **Constant pain** – **gastric carcinoma**
- ❖ **pain attacks** – **acute gastritis, biliary colic**
- ❖ **periodic pain** – **reflux esophagitis** (occurs in the night time and in clinostatism)
- ❖ **hunger/nocturnal pains** (awakes the patient from a deep sleep) - in **duodenal ulcer**
- ❖ **Seasonal pain** - in **peptic ulcer** (exacerbation in spring and autumn)

# Relationship between pain and food intake

- **early postprandial** pain (immediately after ingestion up to 60-90 min postprandial) – reflects an oesophageal or gastric disorder
- **late postprandial** pain (appears in 2-3-4 hours after ingestion; “hunger pain”) - in duodenal ulcer, duodenitis, pancreatic insufficiency

# Antacids and pain relief

---

- Pain in **ulcer** calms down after ingestion of milk, alkaline substances, **H2-blockers**
- Pain in **gastric cancer** does not respond to antacids, but to **opioid analgesics**.

# Extra-Abdominal Causes of the Abdominal pain

## II- Abdominal wall

- 1- Myositis (Bornholm's disease)
- 2- Trauma to abdominal pain
- 3- Muscle strain in cough

## III- Chest causes (referred along intercostal nerves)

- 1- Diaphragmatic pleurisy
- 2- Pneumonia
- 3- Pneumothorax

## IV- CVS causes

- 1- Angina
- 2- MI
- 3- Pericarditis
- 4- CHF

## V- Metabolic and endocrinal causes

- 1- DKA
- 2- Thyrotoxic crisis
- 3- Addisonian crisis
- 4- Acute porphyria
- 5- Severe hypercalcemia

## VI- Neurological causes

- 1- Herpes zoster of lower intercostal nerves
- 2- Referred pain from spinal arthritis

## VII- Other causes

- 1- Torsion of testis



# Abdominal

---



- **Perforation of a gastro-duodenal ulcer**
- **Dissection of aorta,**
- **Rupture of oesophagus,**
- **Extrauterine pregnancy,**
- **Renal stones**

# Dysphagia

---

**Feeling of “blockage” or obstruction of food passage through pharynx or oesophagus, difficulty in swallowing**

# Types of dysphagia

Mechanical dysphagia (organic) –  
caused by **narrowing or intrinsic compression of oesophageal lumen** (carcinoma, post ulcerative strictures, a huge amount - bolus - of food)

**Motorial Dysphagia (functional) –**  
derangements of nervous system or  
musculature, it results in intermittent of  
dysphagia, it comes and goes  
(pharyngeal paralysis, achalasia,  
spasm).

# Mechanical vs Functional dysphagia:

---

1. Difficulty in swallowing of a solid alimentary bolus, and only in advanced stages – including liquid food.
2. Inefficiency of spasmolytics.

1. Difficulty in swallowing of a liquid alimentary bolus, the solid one passes easier.
2. Spasmolytics are efficient.

# Deranged appetite

---

- 1. Increased appetite – duodenal ulcer**
- 2. Anorexia – diminished up to loss of appetite (gastric ulcer, cancer)**
- 3. Bulimia – exaggerated feeling of hunger**
- 4. Aversion for meat – gastric cancer**

**5. Perverse appetite – wish to eat non-edible substances – chalk, soil, newspapers etc.(anaemia, in pregnancy)**

**6. Citofobia – fear of eating (gastric ulcer)**

# Other functional symptoms

---

1. **Aphagia** – complete oesophageal obstruction
2. **Odinophagia** – painful deglutition
3. **Phagophobia** (fear of swallowing and *refuse to swallow*) – in isteria, rabies, tetanus



# Vomiting (or emesis)

- 1. Peripheral– visceral etiology (chronic gastritis, peptic ulcer, pylorostenosis, alcohol abuse)**
- 2. Reflective (outside the stomach) – inferior AMI, appendicitis, peritonitis**
- 3. Central- (vomiting center) – cerebral edema, intracranial pressure, tumors, fever**
- 4. Psychogenic (emotional distress)**
- 5. Hematogenic (toxic) - renal failure, food poisoning, infections, drugs side effects**

# The Act Of Vomiting

**Higher centres:**  
hypothalamus, cerebellum,  
labyrinth, area postrema


**Vomiting centre**

Cascade of vomit



Contraction of diaphragm and abdominal muscles

LES relaxation




Respiration stop

Abdominal pressure

Somatomotor signals

Fig. 22-4

# Mechanism of vomiting



Source: Katzung BG, Masters SB, Trevor AJ: *Basic & Clinical Pharmacology*, 11th Edition: <http://www.accessmedicine.com>

Copyright © The McGraw-Hill Companies, Inc. All rights reserved.

# Complications of vomiting:

- Rupture of oesophagus (Boerhaave symptom)
- Linear ruptures of mucosa in the region of cardio-oesophageal junction (Mallory-Weiss syndrome)
- Dehydration
- Loss of gastric acidity (HCl) – metabolic alkalosis with hypopotasiemia (arrhythmia)

# Heartburn (pyrosis)

- ❑ Burning sensation, retrosternal or in epigastria; irradiates to the neck, sometimes to arms
- ❑ More often is associated with **gastrooesophageal reflux** due to mucosal irritation
- ❑ Relieved by antacids

## **Eructation**

- the passage of gas from the stomach or esophagus through the mouth

## **Regurgitation**

- is the spitting up of food from the esophagus or stomach without nausea or forceful contractions of the abdominal muscles

## **Rumination**

- is regurgitation with no apparent physical cause ( infants, emotional disorders)

# Meteorism

- An increased formation of intraintestinal gas with abdominal distension and flatulence.

## *Appears after:*

- Ingestion of specific aliments (vegetables, some cereals)
- bacterial colonisation of small intestine (*Lambliosis*)



# Diarrhoea

**Increased daily amount of stools over 300g; usually associated with increased fluidity and frequency of stools.**

diarrhoea is considered ***chronic*** after 2 weeks





# Forms of diarrhoea

---

- 1) **Inflammatory**
- 2) **Osmotic**
- 3) **Secretory**
- 4) **Motility disturbances**

# *1. Inflammatory diarrhoea*

- ❖ Parasite infections – helminth, amoeba
- ❖ Infections – salmonella, shigella, E.coli
- ❖ Ulcerative colitis, Crohn disease (autoimmune mechanisms)
- ❖ Colitis due to physical agents: toxins – Hg, Ar, irradiation
- ❖ Ischemic colitis, vasculitis

## 2. *Osmotic diarrhoea*

### **Ingestion of osmotically active products:**

- ❖ **Laxatives**
- ❖ **Products containing sorbitol, xilitol: chewing gum**
- ❖ **Medications: lactulose, almagel (Mg)**

### **Absorption deficiencies**

- ❖ **Deficiency of: disaharide (lactase, sucrose), enterochinase**
- ❖ **Congenital malabsorption**
- ❖ **Exocrine pancreatic insufficiency**
- ❖ **Diminished absorption surface (short intestine, inflammation)**

## *3. Secretory diarrhoea*

- ❖ Infections (cholera, Staphylococcus aureus, Escherichia coli)
- ❖ Tumours
- ❖ Some laxatives
- ❖ Dihydroxilated biliary acids

## *4. Motility disturbances*

### **Hypermotility**

- ❖ Irritable intestine syndrome
- ❖ Carcinoid syndrome (serotonin)
- ❖ Hyperthyroidism

### **Hypomotility**

- ❖ Diabetes mellitus
- ❖ Hypothyroidism
- ❖ Scleroderma
- ❖ Amiloidosis

# Constipation

- stools are less than 3 times/week (1 time in 48 hours).
- secondary to this there is an increased absorption of water – the stool becomes more consistent.
- constipation is considered *chronic* after 6 weeks.

# Causes of constipation

- 1. Colon tumour or foreign body, strictures of the colon, infections, ischemic colitis**
- 2. Psychogenic**
- 3. Functional (reduced intake of liquids, fibres; reduced exercise)**

- 4. Rectal diseases, anal channel diseases**
- 5. Nervous system lesions**
- 6. Metabolic and endocrine diseases**
- 7. Intoxications**
- 8. Digestive system diseases**
- 9. Drugs (analgesic, opiates, antidepressive, antipsychotic, calcium channels blockers)**



# **Gastrointestinal haemorrhage**

**is an emergency, always having an organic reason**

# Signs of GI haemorrhage:

## 1. Haematemesis – vomiting with blood.

if haematemesis happens in short time after onset of bleeding, vomiting masses are **red**.

if haematemesis happens in 0,5 -1 hours, vomiting masses are **dark red, brown or black**, like “**coffee ground**” (blood degraded by HCl)

**2. Melena** – elimination of black stools, like pitch, “like fuel oil”, caused by blood from an upper gastrointestinal haemorrhage (oesophagus, stomach or duodenum), digested by microbial flora and becoming dark

*Lesions of jejunum, ileum and ascending colon* can cause **melena**, when the time of gastrointestinal transit is prolonged

---

**3. Haematochezia - passage of red blood through rectum, as a sign of bleeding from a distal source (Treitz ligament).**

# Severity of haemorrhage:

< 500 ml – without clinical signs

signs of hypovolemic shock (loss of more than 40% blood volume):

**Lipothymia, syncope, nausea, transpiration and thirst**

**Pale and cold skin**

**Agitation**

**Arterial hypotension**

**Tachycardia**

## ***Aetiology of superior DH***

- erosive or hemorrhagic gastropathy (NAID, anticoagulants, alcohol),
- duodenal or gastric ulcer,
- s-m Mallory-Weiss,
- oesophageal varices
- malign tumours
- oesophagitis (5-8%),
- duodenitis (5-9%),
- angiodysplasia (5-7%),

## *Aetiology of inferior DH*


- **anorectal disease**


---
- **polyps, cancer**
- **diverticulosis**
- **abnormal intestinal tract**
- **enterocolitis, colitis, intestinal ischemia**

# Inspection


- The patient relaxed and comfortable in supine position
- Use relaxation techniques if needed Head supported with pillow
- Keep the supinated arm by patient sides, warm hands
- Insure good illumination, full exposure of the abdomen
- Be on the right side of the patient







# Normal



# Inspection of abdomen

Contours and peristalsis of the abdomen

Abdomen dimensions

Abdomen symmetry

Diastasis recti


Presence of local bulging (hernia, tumor)

Skin and subcutaneous fat

- Umbilicus inspection (Position and protrusion)

Superficial venous circulation (caput Medusae )

# Common abdominal scars




# Obesity



# Hepatomegaly



# Gallbladder, Jaundice





# Ascites, protrusion of umbilicus



# Umbilicalal hernia



# Superficial venous circulation



**A**

**Cullen's sign**




**B**

**Grey Turner's sign**



# Signs of CLD



# Auscultation I

Provides important information about bowel motility:

a. decreased motility suggests peritonitis

b. increased motility suggests obstruction


- *Need to listen before percussion or palpation since these maneuvers may alter the frequency of bowel sounds*

- Can also appreciate bruits over the aorta or other arteries, suggesting narrowing of the arteries from atherosclerosis

# Auscultation II



- 
- Listen with diaphragm of stethoscope
  - Normal sounds occurs every 5-10 seconds & consist of clicks and gurgles
  - Need to listen for 2 minutes to declare no bowel sounds; since bowel sounds are widely transmitted, need only to listen in one spot
  - Occasionally hear *borborygmi* - long, prolonged gurgles of hyperperistalsis - the familiar stomach growling








Fig. 17-8 Sites to auscultate for bruits: renal arteries, iliac arteries, aorta, and femoral arteries.


# Percussion

---

Helps to identify the amount and distribution of gas and to identify possible masses that are solid or fluid filled

- Can be used to assess size of liver and spleen
- Percuss looking for areas of tympany and dullness
- Large dull areas may indicate an underlying mass; you will later confirm with palpation
- On the right is liver dullness; on the left, dullness of the spleen

# Assessment of liquid in abdominal cavity



# Assessment of liquid in abdominal cavity

—

## fluctuation (wave) sign



# Percussion



# Palpation

superficial (light)

deep

# Light and Deep Palpation

---

- Light palpation
  - Helpful in identifying tenderness, superficial organs, masses, hernia of medial abdominal line, Blumberg symptom
  - Palpate with a light, gentle dipping motion using the palmar surface of fingers
- Deep palpation
  - Usually required to delineate abdominal masses
  - Again use palmar surface of fingers
  - Check for tenderness and rebound (pain induced or increased by letting go)

# Palpation: Improving the Exam

---

Patient should have an empty bladder

- Patient supine, arms at sides or folded across chest
  - avoid arms above the head as this tightens the abdomen
- Before you begin, ask the patient to point to areas of pain and examine last
- Warm hands and stethoscope; avoid long nails; approach slowly
- Distract the patient with conversation or questions



# General rules of palpation

---

1. The doctor is sitting on the right part of the patient, at the level of the bed
2. The painful region of the abdomen is to be palpated at the end

# The order of *superficial palpation* (*counterclockwise*):

---

**Left inguinal region**

**Left flank (Left lateral region)**

**Left hypochondria**

**Epigastria**


**Right hypochondria**

**Right flank**

**Right inguinal region**

**Suprapubian region**

**Umbilical region**



# Blumberg symptom

**sign of irritation of peritoneum (positive in peritonitis).**

## **Appreciation:**

- 1. The palpatory hand pushes the abdomen in the painful region (pain is present),**
- 2. Take off abruptly the hand. If pain is intensified the Blumberg sign is positive.**

# Deep palpation

**Aim – direct examination of different parts of gastrointestinal tract.**

## Appreciate:

**Dimensions**

**Shape**

**Presence of irregularities**

Several structures are palpable normally:

- Sigmoid colon is frequently palpable as a firm, narrow tube in the left lower quadrant

---

- The caecum and ascending colon form a softer, wider tube in the right lower quadrant

- Normal liver distends below the costal margin but its soft consistency is difficult to feel

- Pulsations of the abdominal aorta are frequently visible and usually palpable

- **Usually NOT palpable are: stomach, spleen, gallbladder, duodenum, pancreas, kidneys**

# The order of *deep palpation*:

---

- 1. Sigmoid colon**
- 2. Caecum**
- 3. Terminal segment of ileum**
- 4. Ascending colon**
- 5. Descending colon**
- 6. Transversal colon**
- 7. Big curvature of the stomach**
- 8. Pylorus**

# Method of deep palpation of the abdomen

---

There are 4 consecutive steps



1 moment – apply the hand parallel to the palpated margin; the other hand is on the abdomen in order to calm down the patient



## 2 moment – form the skin folder

---



# The skin folder is to be formed

---

## To the umbilicus


For part of the colon, situated below the umbilicus (sigmoid colon, caecum, ileocaecal angle, ascending colon, descendent colon)

## From the umbilicus

For parts, situated above the umbilicus (transversal colon, stomach)


3 moment – deepening the fingers into abdomen (in expiration)

---



4 moment – sliding on the surface of the respective organ

---



# Palpation of ascending and descending colon

---



# Palpation of ascending and descending colon

---

The left hand is on the posterior part of the abdomen (in lumbar region), moving the tissues to the hand which is doing palpation (right)

# Palpation of the big curvature of the stomach

---





# Paraclinical examination

BARIUM SWALLOW

UPPER GI ENDOSCOPY

ENDOSCOPIC RETROGRADE CHOLANGIOPANCREATOGRAPHY  
(ERCP)

PANCREAS SCAN

LIVER SCAN

LIVER BIOPSY

COLONOSCOPY




SIGMOIDOSCOPY

ABDOMINAL X-RAY

ABDOMINAL ULTRASOUND

CT SCAN OF THE ABDOMEN

LAPAROSCOPY



Level of cut section



